

## ACKNOWLEDGEMENTS

I would like to convey heartfelt gratitude to the following people and institutions:

- The University of Zimbabwe-Flemish Universities Link (UZ-VLIR Training of Fisheries Biologists Project) for generously funding this project
- My supervisor, Prof. B.E. Marshall, as well as Dr B.N. Dube (UZ), Professors F. Ollevier and L. Brendonck (KU Leuven, Belgium) for their constructive criticism
- Academic and technical staff of the UZ Biological Sciences Department for their support, particularly Ms C. Mpamhanga, Mr S. Alferi, Mr G. Ashley, Mr E. Mpomba and the late Mr Kanengoni, Mr R. Nhavira and Mr Tambanemoto (may their souls rest in peace)
- Fellow students Tamuka Nhiwatiwa, Brian Gratwicke, Tinashe Muteveri, Zviregei Mbizi, Nzwirashe Dekeza & Chenjerai Kashangura for field assistance
- The research assistants T. Zengeya, C. Kanonda, R. Mombeshora, E. Mamire, W. Chipamuriwo, H. Munemo, B. Gweshe, Dennis and T. Kunze for fieldwork assistance
- Prof. N.M. Mokgalong, Mrs W. Luus-Powell, Mrs A. Moller and Mr M. Matla (University of the North, South Africa) for assisting in identifying trematode and nematode parasites from fish and birds, and for technical assistance during scanning electron microscopy work
- Prof. S.N. Mashego (University of the North-West, South Africa) for assisting in the identification of fish cestodes and nematodes
- Prof. A. Avenant-Oldewage (Rand Afrikaans University, South Africa) for literature support
- The Department of National Parks and Wildlife Management for granting sampling permits
- Mr C. Phiri (formerly senior Ecologist – Aquatic) and Mrs T. Couto (Senior Ranger – Ornithology, Lake Chivero Fisheries Research Station) for field assistance
- Mr D. Stewart of Ingwerati Farm, for permission to sample in the Munwahuku Dam
- Prof. Mukaratirwa (Dean) and Mr Taruvinga (Senior Technologist – Department of Paraclinical Veterinary Sciences, UZ) for technical assistance (laboratory methods)
- My family and friends for their spiritual and moral support
- Finally, but not least, to my dear wife Shupikai, “...*behind every successful man is a successful, beautiful, hardworking wife*”, and to my son Valentine Tadiwanashe, “...*like father, like son*”.

**MAY GOD BLESS YOU ALL!**

## CONTENTS

	Page
Acknowledgements	i
List of figures	iv
List of tables	vi
Chapter 1. General Introduction	1
Chapter 2. The occurrence of <i>Contraecaecum</i> species (Nematoda: Anisakidae) in the sharptooth catfish, <i>Clarias gariepinus</i> , and fish-eating birds from Lake Chivero, Zimbabwe	10
2.1. Introduction	10
2.2. Materials and Methods	12
2.3. Results	15
2.4. Discussion	21
Chapter 3. Tapeworms of fish: The occurrence of <i>Ligula intestinalis</i> in <i>Barbus paludinosus</i> and <i>Proteocephalus</i> sp. in <i>Clarias gariepinus</i>	27
3.1. Introduction	27
3.2. Materials and Methods	29
3.3. Results	31
3.4. Discussion	38
Chapter 4. Trematode parasites of fish: The occurrence of <i>Harvardia sandgroundi</i> (Diplostomidae) metacercariae in <i>Barbus paludinosus</i> from the Marimba River and <i>Clinostomum complanatum</i> (Clinostomidae) in piscivorous birds from Lake Chivero	45
4.1. Introduction	45
4.2. Materials and Methods	47
4.3. Results	49
4.4. Discussion	50
Chapter 5. Taxonomic notes on the major helminth species of <i>Barbus paludinosus</i> , <i>Clarias gariepinus</i> and piscivorous birds from the Manyame catchment area	52
Chapter 6. Conclusions	73
References	76

## FIGURES

		Page
1.1.	Location of lake Chivero, Munwahuku Dam and the Marimba River in relation to the city of Harare	9
2.1.	The relation between parasite load (intensity) and the condition factor (K) of <i>C. gariepinus</i> from Lake Chivero	16
2.2.	The variation in (a) prevalence and (b) mean intensity of <i>Contracaecum</i> infection of <i>C. gariepinus</i> of different size classes and sexes	18
3.1.	The monthly prevalence of <i>Ligula intestinalis</i> in <i>Barbus paludinosus</i> from the Munwahuku Dam (vertical bars), and the abundance of copepods (calanoids + cyclopoids) collected from the dam (line plot)	34
3.2.	The seasonal prevalence of <i>Ligula intestinalis</i> in <i>Barbus paludinosus</i> from the Munwahuku Dam. Each season is represented by the median month	35
3.3.	The relationship between the weight of the host ( <i>B. paludinosus</i> ) and the weight of the parasite ( <i>L. intestinalis</i> ) from the Munwahuku Dam. Dotted lines indicate the 95% confidence interval range	36
3.4.	The relationship between the weight of <i>Ligula intestinalis</i> and the condition factor (K) of <i>Barbus paludinosus</i> from the Munwahuku Dam	37
3.5.	The parasite load per fish (intensity) of <i>B. paludinosus</i> from the Munwahuku Dam and that of four other <i>Barbus</i> species from the Middle Letaba Dam (Saayman <i>et al.</i> , 1991) expressed as a percentage of the total number (N) of infected hosts	40
5.1.	<i>Harvardia sandgroundi</i> – photomicrographs of (a) forebody and (b) hindbody (scale bar = 0.8mm). (c) Drawing to illustrate features in (a) and (b) (Scale bar = 1.0mm).	55
5.2.	<i>Clinostomum complanatum</i> from the buccal cavity of <i>Phalacrocorax carbo</i> (Lake Chivero). Scale bar = 1.0mm	57
5.3.	Photomicrographs of <i>Proteocephalus</i> sp. (a) scolex, (b) mature proglottide showing arrangement of testes and vitellaria, (c) mature proglottide showing position of ovaries and uterus, (d) scanning electron micrograph of scolex of an immature adult. Note that the suckers are not yet fully developed.	60

- 5.4. The head of a female *Contracaecum rudolphii* from the Reed Cormorant showing two lobes in interior margin of each lip. Scale bar = 10 $\mu$ m 66
- 5.5. The tail of a female *C. rudolphii* from the White-breasted Cormorant. Note the absence of pre- and postcloacal papillae. Scale bar = 100 $\mu$ m 66
- 5.6. The tail of a male *C. rudolphii* from the White-breasted Cormorant showing the arrangement of postcloacal papillae. Note the two double papillae, lateroventral and midway between tip of tail and cloacal opening. Scale bar = 10 $\mu$ m 67
- 5.7. The tail of a male *C. rudolphii* from the Reed Cormorant showing the spicules (fused) and papillae. Scale bar = 100 $\mu$ m 67
- 5.8. The head of *C. carlislei* from the White-breasted Cormorant characterized by somewhat rectangular lips and two single papillae on dorsal lip. Scale bar = 100 $\mu$ m 68
- 5.9. The tail of a male *C. carlislei* from the White-breasted Cormorant. Note the subequal spicules. Scale bar = 100 $\mu$ m 68
- 5.10. The tail of a female *C. carlislei* from the White-breasted Cormorant. Scale bar = 100 $\mu$ m 69
- 5.11. The head of *C. microcephalum* from the Darter. Note the hexagonal lips. Scale bar = 100 $\mu$ m 69
- 5.12. The tail of a male *C. microcephalum* from the Darter with subequal spicules. Note the longitudinal, wrinkle-like alae. Scale bar = 100 $\mu$ m 70
- 5.13. The tail of a male *C. microcephalum* from the Darter showing the arrangement of papillae. Scale bar = 10 $\mu$ m 70
- 5.14. The head of *C. tricuspis* from the Darter showing the characteristic tri-forked interlabia. Scale bar = 100 $\mu$ m 71
- 5.15. The tail of a male *C. tricuspis* from the Darter showing papillae and subequal spicules. Note the absence of lateral alae. Scale bar = 100 $\mu$ m 71
- 5.16. The tail of *C. tricuspis* from the Darter showing the arrangement of papillae. Scale bar = 100 $\mu$ m 72

## TABLES

	Page
1.1. Some publications on freshwater fish parasites from African countries	5
1.2. Some physical and chemical characteristics of Lake Chivero (from Elenbaas and Grundel, 1994), Munwahuku Dam and Marimba River	8
2.1. The prevalence (% infected) of <i>Contracaecum</i> larvae in <i>C. gariepinus</i> from Lake Chivero (November 2000 – May 2002). N = number of fish sampled	17
2.2. The infection dynamics of <i>Contracaecum</i> species found in the alimentary canal of four species of piscivorous birds at Lake Chivero. MI = mean intensity	19
2.3. The diet of four avian species collected from Lake Chivero. Pa = <i>Pharyngochromis acuticeps</i> , On = <i>Oreochromis niloticus</i> , Cc = <i>Cyprinus carpio</i> , uc = unidentified cichlids, ND = not determined	20
2.4. The parasitic load of <i>Contracaecum</i> larvae in <i>C. gariepinus</i> from Lake Chivero in comparison with other dams in Africa	23
3.1. The occurrence and prevalence of cestode parasites in <i>Barbus paludinosus</i> and <i>Clarias gariepinus</i> from Munwahuku Dam, Lake Chivero and the Marimba River. P = prevalence (%), I = intensity, MI = mean intensity	32
3.2. The mean abundance (numbers x 10 <sup>3</sup> m <sup>-3</sup> ) of copepods (calanoids + cyclopoids) collected from the Munwahuku Dam, Lake Chivero and the Marimba River and the Munwahuku Dam	32
3.3. The prevalence and intensity of <i>Ligula</i> parasitism in <i>B. paludinosus</i> from the Munwahuku Dam compared with some <i>Barbus</i> and <i>Labeobarbus</i> species from South African impoundments. N = sample size, P = prevalence, I = intensity, MI = mean intensity. Sources are (1) Saayman <i>et al.</i> (1991); (2) Mashego (1982); (3) this study.	39
3.4. A comparison of the parasite index (PI) values of <i>B. paludinosus</i> collected from Munwahuku Dam with those of other <i>Barbus</i> species from Middle Letaba Impoundment, South Africa (from Saayman <i>et al.</i> , 1991).	44
3.5. The prevalence and intensity of <i>Proteocephalus</i> infection in <i>C. gariepinus</i> from some dams in South Africa compared with Lake Chivero. N = sample size, P = prevalence, I = intensity, MI = mean intensity.	44

**TABLES (cont.)**

	Page
4.1. Diplostomid metacercariae on the skin of <i>Barbus paludinosus</i> from the Marimba River. N = sample size, P = prevalence (%), I = intensity, MI = mean intensity.	49
5.1. Measurements (millimetres) of <i>Harvardia sandgroundi</i> metacercariae from <i>Barbus paludinosus</i> collected in the Marimba River	54
5.2. Measurements (micrometers) of <i>Clinostomum complanatum</i> from birds on Lake Chivero compared with those recorded by other workers. Mean values in parentheses.	56
5.3. Morphological features used in identification of L3 larvae of <i>Contracaecum</i> sp. in <i>Clarias gariepinus</i> from Lake Chivero. All measurements are in mm.	62